

BC

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
31 July 2003 (31.07.2003)

PCT

(10) International Publication Number
WO 03/062190 A1(51) International Patent Classification⁷: C07C 255/41,
309/73, A61K 31/165[CA/CA]; 10 Ruddington Drive, Apt 303, Toronto, Ontario
M2K 2J7 (CA).

(21) International Application Number: PCT/CA03/00032

(74) Agent: LEDGLEY, Cynthia, J.; 121 Richmond Street
West, Suite 1000, Toronto, Ontario M5H 2K1 (CA).

(22) International Filing Date: 17 January 2003 (17.01.2003)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE,
SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VC, VN, YU, ZA, ZM, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/349,215	18 January 2002 (18.01.2002)	US
60/349,214	18 January 2002 (18.01.2002)	US
60/349,216	18 January 2002 (18.01.2002)	US

(71) Applicant (*for all designated States except US*): THE
HOSPITAL FOR SICK CHILDREN [CA/CA]; 555
University Avenue, Suite 5270, Toronto, Ontario M5G
1X8 (CA).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): ROIFMAN, Chaim,
M. [CA/CA]; 33 Christine Crescent, Toronto, Ontario
M2R 1A4 (CA). DEMIN, Peter [CA/CA]; 414-40
High Park Avenue, Toronto, Ontario M6P 2S1 (CA).
GRUNBERGER, Thomas [CA/CA]; 63 Robingrove
Road, Toronto, Ontario M2R 3A1 (CA). ROUNOVA,
Olga [RU/CA]; 414-40 High Park Avenue, Toronto,
Ontario M6P 2S1 (CA). CIMPEAN, Octavian, Laurand

Declaration under Rule 4.17:



— of inventorship (Rule 4.17(iv)) for US only

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: COMPOUNDS FOR MODULATING CELL PROLIFERATION

(57) Abstract: The following compounds of formulae (I) and (III) which may
be useful in treating a variety of cell proliferative disorders are disclosed: Fo-
mula (I), (II), wherein R¹, R² and R³ are each independently selected from H,
OH, C₁₋₆alkyl, C₁₋₆alkoxy, NH₂, NH-C₁₋₆alkyl, N(C₁₋₆alkyl)(C₁₋₆alkyl), SH,
S-C₁₋₆alkyl, NO₂, CF₃, OCF₃ and halo; R⁴ is unsubstituted Ar, or Ar substi-
tuted with 1-4 substituents, independently selected from C₁₋₆alkyl, C₁₋₆alkoxy
and halo; X is selected from (CH₂CH₂O)_n and (CH₂)_n, and n=1-4.

BEST AVAILABLE COPY

WO 03/062190 A1

COMPOUNDS FOR MODULATING CELL PROLIFERATION

Field of the Invention

This invention relates to compounds and their use for treating a variety of cell proliferative disorders.


Background of the Invention

A number of small molecules which act as tyrosine kinase inhibitors have been identified. For example, certain compounds have been described as tyrosine kinase inhibitors, effective to inhibit cell proliferation (see for example, US 5,891,917, US 5,217,999, US 5,773,476, US 5,935,993, US 5,656,655, US 5,677,329, US 5,789,427 and WO 01/79158). However, there remains a need for compounds that selectively target one or more such kinases, e.g., to preferentially inhibit proliferation of cancerous cells over normal cell proliferation.

Summary of the Invention

Compounds are provided that may be used for modulating abnormal cell proliferation, for example inhibiting cancer cell proliferation, and which preferably do not adversely affect normal cell proliferation.

Accordingly, the present invention includes compounds of Formula I and salts, solvates and hydrates thereof:



wherein


R^1 , R^2 , R^3 are each independently selected from H, OH, C $_{1-6}$ alkyl, C $_{1-6}$ alkoxy, NH $_2$, NH-C $_{1-6}$ alkyl, N(C $_{1-6}$ alkyl)(C $_{1-6}$ alkyl), SH, S-C $_{1-6}$ alkyl, NO $_2$, CF $_3$, OCF $_3$ and halo;

R^4 is unsubstituted Ar, or Ar substituted with 1-4 substituents, independently selected from C₁₋₆alkyl, C₁₋₆alkoxy and halo;

X is selected from (CH₂CH₂O)_n and (CH₂)_n, and
n = 1-4.

5 In certain embodiments, X is (CH₂)_n.

The present invention also provides compounds of Formula II and salts, solvates and hydrates thereof:




10 wherein

R^1 , R^2 , R^3 are each independently selected from H, OH, C₁₋₆alkyl, C₁₋₆alkoxy, NH₂, NH-C₁₋₆alkyl, N(C₁₋₆alkyl)(C₁₋₆alkyl), SH, S-C₁₋₆alkyl, NO₂, CF₃, OCF₃ and halo;

R^4 = C₁₋₆alkyl; and

15 n = 1-4.

Additionally, the present invention includes compounds of Formula III and salts, solvates and hydrates thereof:



20

wherein

R^1 , R^2 and R^3 are each independently selected from H, OH, C₁₋₆alkyl, C₁₋₆alkoxy, NH₂, NH-C₁₋₆alkyl, N(C₁₋₆alkyl)(C₁₋₆alkyl), SH, S-C₁₋₆alkyl, NO₂, CF₃, OCF₃ and halo; and

25


R^4 is selected from C₁₋₆alkyl, phenyl and pyridyl, wherein phenyl and pyridyl are unsubstituted or substituted with 1-4 substituents, independently selected

3

from C₁₋₆alkyl, C₁₋₆alkoxy and halo, with the provisos that when R¹ and R³ are both H and R⁴ is unsubstituted phenyl, R² is not H, Cl, or OCH₃; when R¹ and R² are both H and R⁴ is unsubstituted phenyl, R³ is not NO₂; and when R¹ and R³ are both H and R⁴ is CH₃, R² is not N(CH₃)₂.

5

In accordance with a further aspect of the present invention, there are provided compounds of Formula IV and salts, solvates and hydrates thereof, for use in compositions and methods for treating cell proliferation:



IV

10

wherein

R¹, R² and R³ are each independently selected from H, OH, C₁₋₆alkyl, C₁₋₆alkoxy, NH₂, NH-C₁₋₆alkyl, N(C₁₋₆alkyl)(C₁₋₆alkyl), SH,

15 S-C₁₋₆alkyl, NO₂, CF₃, OCF₃ and halo; and

R⁴ is selected from C₁₋₆alkyl, phenyl and pyridyl, wherein phenyl and pyridyl are unsubstituted or substituted with 1-4 substituents, independently selected from C₁₋₆alkyl, C₁₋₆alkoxy and halo.

20

According to another aspect of the present invention, there is provided a pharmaceutical composition comprising a compound as set forth herein and a pharmaceutically acceptable carrier or diluent.

25 In accordance with a further aspect of the present invention, there is provided a method for modulating cell proliferation, preferably inhibiting cell proliferation, comprising administering an effective amount of a compound as set forth herein to a cell or animal in need thereof. The invention also includes a use of a compound as set forth herein to modulate cell proliferation, preferably inhibit cell proliferation, for example in cancer cells. The invention further provides for a use of a compound

as set forth herein to prepare a medicament to modulate cell proliferation, preferably inhibit cell proliferation.

In another aspect, the invention provides a method of modulating tyrosine kinase activity in a cell by administering an effective amount of a compound as set forth herein. In a further aspect, the invention provides a method of inhibiting tyrosine kinase activity in a cell by administering an effective amount of a compound as set forth herein. The present invention also provides a use of a compound as set forth herein to modulate, preferably inhibit, tyrosine kinase activity. The present invention further provides a use of a compound as set forth herein to prepare a medicament to modulate tyrosine kinase activity, preferably inhibit tyrosine kinase activity. It is appreciated that the modulation of cell proliferation by the compounds herein may be effected by other mechanisms and that tyrosine kinase modulation may be just one mechanism. Accordingly, the compounds described herein are useful in the compositions and methods set forth herein without regard to the actual mechanism contributing to their effectiveness.

Other features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples while indicating preferred embodiments of the invention are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

25 Detailed Description of the Preferred Embodiments

I. Definitions

The term "C₁₋₆alkyl" as used herein means, unless otherwise stated, straight and/or branched chain alkyl radicals containing from one to six carbon atoms and includes methyl, ethyl, propyl, isopropyl, t-butyl and the like.

5

The term "C₁₋₆alkoxy" as used herein means, unless otherwise stated, straight and/or branched chain alkoxy radicals containing from one to six carbon atoms and includes methoxy, ethoxy, propyloxy, isopropyloxy, t-butoxy and the like.

5 The term "C₁₋₄alkyl" as used herein means, unless otherwise stated, straight and/or branched chain alkyl radicals containing from one to four carbon atoms and includes methyl, ethyl, propyl, isopropyl, t-butyl and the like.

The term "C₁₋₄alkoxy" as used herein means, unless otherwise stated, straight
10 and/or branched chain alkoxy radicals containing from one to four carbon atoms and includes methoxy, ethoxy, propyloxy, isopropyloxy, t-butoxy and the like.

The term "halo" as used herein means halogen and includes chloro, fluoro, bromo, iodo and the like.

15

The term "pharmaceutically acceptable salt" means an acid addition salt which is suitable for or compatible with the treatment of patients.

The term "compound(s) as set forth herein" as used herein includes any
20 compound of the Formulae I, II, III, and IV as defined herein (including all salts, solvates, or hydrates thereof) as well as any specific compound designated herein, such as CRIX-38, CRIX-39, CRIV-42, CRIV-46, CRVIII-33, CRVIII-34, CRVIII-35, CRVIII-50, CRVIII-51, CRVIII-52, CRVIII-53, and CRIX-79 (including all salts, solvates or hydrates thereof).

25

The term "pharmaceutically acceptable acid addition salt" as used herein means any non-toxic organic or inorganic salt of any base compounds represented by Formulae I-IV, or their intermediates. Illustrative inorganic acids which form suitable salts include hydrochloric, hydrobromic, sulfuric and phosphoric acids, as
30 well as metal salts such as sodium monohydrogen orthophosphate and potassium hydrogen sulfate. Illustrative organic acids that form suitable salts include mono-, di-, and tricarboxylic acids such as glycolic, lactic, pyruvic, malonic, succinic,

glutaric, fumaric, malic, tartaric, citric, ascorbic, maleic, benzoic, phenylacetic, cinnamic and salicylic acids, as well as sulfonic acids such as *p*-toluenesulfonic and methanesulfonic acids. Either the mono or di-acid salts can be formed, and such salts may exist in either a hydrated, solvated or substantially anhydrous form. In general, the acid addition salts of compounds as set forth herein are more soluble in water and various hydrophilic organic solvents, and generally demonstrate higher melting points in comparison to their free base forms. The selection of the appropriate salt will be known to one skilled in the art. Other non-pharmaceutically acceptable salts, e.g. oxalates, may be used, for example, in the isolation of compounds of Formulae I-IV for laboratory use, or for subsequent conversion to a pharmaceutically acceptable acid addition salt.

The term "solvate" as used herein means a compound of Formulae I-IV, or a pharmaceutically acceptable salt of a compound of Formulae I-IV, wherein molecules of a suitable solvent are incorporated in the crystal lattice. A suitable solvent is physiologically tolerable at the dosage administered. Examples of suitable solvents are ethanol, water and the like. When water is the solvent, the molecule is referred to as a "hydrate".

The term an "effective amount" or a "sufficient amount" of an agent as used herein is that amount sufficient to effect beneficial or desired results, including clinical results, and, as such, an "effective amount" depends upon the context in which it is being applied and would be understood by a person skilled in the art. For example, in the context of administering an agent that modulates cell proliferation, an effective amount of an agent is, for example, an amount sufficient to achieve such a modulation in cell proliferation as compared to the response obtained without administration of the agent.

As used herein, and as well understood in the art, "treatment" is an approach for obtaining beneficial or desired results, including clinical results. Beneficial or desired clinical results can include, but are not limited to, alleviation or amelioration of one or more symptoms or conditions, diminishment of extent of disease,

7

stabilized (i.e., not worsening) state of disease, preventing spread of disease, delay or slowing of disease progression, amelioration or palliation of the disease state, and remission (whether partial or total), whether detectable or undetectable. "Treatment" can also mean prolonging survival as compared to expected survival if

5 not receiving treatment.

"Palliating" a disease or disorder means that the extent and/or undesirable clinical manifestations of a disorder or a disease state are lessened and/or time course of the progression is slowed or lengthened, as compared to not treating the

10 disorder.

The term "prevention" is art-recognized, and when used in relation to a condition, such as a local recurrence (e.g., pain), a disease such as cancer, a syndrome complex such as heart failure or any other medical condition, is well

15 understood in the art, and includes administration of a composition which reduces the frequency of, or delays the onset of, symptoms of a medical condition in a subject relative to a subject which does not receive the composition. Thus, prevention of cancer includes, for example, reducing the number of detectable cancerous growths in a population of patients receiving a prophylactic treatment

20 relative to an untreated control population, and/or delaying the appearance of detectable cancerous growths in a treated population versus an untreated control population, e.g., by a statistically and/or clinically significant amount. Prevention of an infection includes, for example, reducing the number of diagnoses of the infection in a treated population versus an untreated control population, and/or

25 delaying the onset of symptoms of the infection in a treated population versus an untreated control population. Prevention of pain includes, for example, reducing the frequency of, or alternatively delaying, pain sensations experienced by subjects in a treated population versus an untreated control population.

30 The term "modulate" as used herein includes the inhibition or suppression of a function or activity (such as cell proliferation) as well as the enhancement of a function or activity.

To "inhibit" or "suppress" or "reduce" a function or activity, such as cancer cell proliferation is to reduce the function or activity when compared to otherwise same conditions except for a condition or parameter of interest, or alternatively, as
5 compared to another conditions.

To "enhance" or "increase" a function or activity, such as cell proliferation, is to increase the function or activity when compared to otherwise same conditions except for a condition or parameter of interest, or alternatively, as compared to
10 another conditions.

The term "animal" as used herein includes all members of the animal kingdom including human. The animal is preferably a human.


15 The term "a cell" as used herein includes a plurality of cells. Administering a compound to a cell includes *in vivo*, *ex vivo* and *in vitro* administration.

The term "abnormal cell" as used herein includes any cell type that is causative or involved in a disease or condition and wherein it is desirable to
20 modulate, enhance or inhibit proliferation of the abnormal cell to treat the disease or condition.

II. Compounds of the Invention

Compounds are provided which may be used to modulate cell proliferation.
25 As such the compounds may be useful in treating cell proliferative diseases such as cancer.

Accordingly, the present invention provides compounds of Formula 1, and salts, solvates or hydrates thereof:



wherein

R^1 , R^2 and R^3 are each independently selected from H, OH, C_{1-6} alkyl, C_{1-6} alkoxy,

NH_2 , $NH-C_{1-6}$ alkyl, $N(C_{1-6}alkyl)(C_{1-6}alkyl)$, SH,

10 $S-C_{1-6}alkyl$, NO_2 , CF_3 , OCF_3 and halo;

R^4 is unsubstituted Ar, or Ar substituted with 1-4 substituents, independently selected from C_{1-6} alkyl, C_{1-6} alkoxy and halo;

X is selected from $(CH_2CH_2O)_n$ and $(CH_2)_n$, and

$n = 1-3$.

15

In embodiments of the invention, compounds of Formula 1 are those in which R^1 , R^2 and R^3 are each independently selected from H, OH, C_{1-6} alkyl, C_{1-6} alkoxy, NH_2 , $NH-C_{1-6}alkyl$, $N(C_{1-6}alkyl)(C_{1-6}alkyl)$, SH, $S-C_{1-6}alkyl$, NO_2 , CF_3 , OCF_3 and halo. In preferred embodiments, R^1 , R^2 and R^3 are each independently selected from H, OH, $C_{1-4}alkyl$, $C_{1-4}alkoxy$, NH_2 , $NH-C_{1-4}alkyl$, $N(C_{1-4}alkyl)(C_{1-4}alkyl)$, NO_2 , CF_3 , OCF_3 and halo. In more preferred embodiments, R^1 , R^2 and R^3 are each independently selected from H, OH, OCH_3 , NH_2 , $N(CH_3)_2$ and NO_2 . In the most preferred embodiments, R^1 , R^2 and R^3 are each independently selected from H, OH and OCH_3 .

25


Further embodiments of the invention include compounds of Formula 1 wherein R^4 is Ar. In preferred embodiments, R^4 is unsubstituted Ar. Most preferably, R^4 is phenyl.

10

Further embodiments of the invention include compounds of Formula I wherein X is $(CH_2)_n$. Preferably n is 1-3; most preferably n is 1.

In a preferred embodiment of the present invention, compounds of Formula I
5 include those in which at least one of R^1 , R^2 and R^3 is OH, more preferably at least
two of R^1 , R^2 and R^3 are OH, while R^4 is Ar and n is 1-3.

The present invention also provides a compound of Formula II and salts, solvates and hydrates thereof:



wherein

R¹, R², R³ are each independently selected from H, OH, C₁₋₆alkyl, C₁₋₆alkoxy, NH₂, NH-C₁₋₆alkyl, N(C₁₋₆alkyl)(C₁₋₆alkyl), SH, S-C₁₋₆alkyl, NO₂, CF₃, OCF₃ and halo;

$R^4 = C_{1-6}alkyl$; and
 $n = 1-4$.

In embodiments of the invention, compounds of Formula II are those in which R¹, R² and R³ are each independently selected from H, OH, C₁₋₆alkyl, C₁₋₆alkoxy, NH₂, NH-C₁₋₆alkyl, N(C₁₋₆alkyl)(C₁₋₆alkyl), SH, S-C₁₋₆alkyl, NO₂, CF₃, OCF₃ and halo. In preferred embodiments, R¹, R² and R³ are each independently selected from H, OH, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, NH-C₁₋₄alkyl, N(C₁₋₄alkyl)(C₁₋₄alkyl), NO₂, CF₃, OCF₃ and halo. In more preferred embodiments, R¹, R² and R³ are each independently selected from H, OH, OCH₃, NH₂, N(CH₃)₂ and NO₂. In the most preferred embodiments, R¹, R² and R³ are each independently selected from H, OH and OCH₃.

11

Further embodiments of the invention include compounds of Formula II wherein R^4 is C_{1-6} alkyl. In preferred embodiments, R^4 is methyl or ethyl. Most preferably, R^4 methyl.

5 Further embodiments of the invention include compounds of Formula II wherein n in 1-4. Preferably, n is 2-3; most preferably, n is 3.

In specific embodiments of the present invention, a compound of the invention is:

10

2-Cyano-5-(4-hydroxy-3,5-dimethoxyphenyl)-penta-2E,4E-dienoic acid benzyl ester (CRIX-38)

2-Cyano-5-(3,4-dihydroxyphenyl)-penta-2E,4E-dienoic acid benzyl ester (CRIX-39)

2-Cyano-5-(3,4-dihydroxyphenyl)-penta-2E,4E-dienoic acid 2-[2-(2-

15


methoxyethoxy)ethoxy] ethyl ester (CRIV-42)

2-Cyano-5-(4-hydroxy-3,5-dimethoxyphenyl)-penta-2E,4E-dienoic acid 2-[2-(2-methoxyethoxy)ethoxy]ethyl ester (CRIV-46); and

2-Cyano-5-(4-hydroxy-3-methoxyphenyl)-penta-2E,4E-dienoic acid benzyl ester (CRIX-79).

20

Accordingly, the present invention provides compounds of Formula III, and salts, solvates or hydrates thereof:



III

25

wherein

R^1 , R^2 and R^3 are each independently selected from H, OH, C_{1-6} alkyl, C_{1-6} alkoxy, NH_2 , $NH-C_{1-6}$ alkyl, $N(C_{1-6}$ alkyl)(C_{1-6} alkyl), SH, $S-C_{1-6}$ alkyl, NO_2 , CF_3 , OCF_3 and halo; and

12

R^4 is selected from C_{1-6} alkyl, phenyl and pyridyl, wherein phenyl and pyridyl are unsubstituted or substituted with 1-4 substituents, independently selected from C_{1-6} alkyl, C_{1-6} alkoxy and halo, with the provisos that when R^1 and R^3 are both H and R^4 is unsubstituted phenyl, R^2 is not H, Cl, or OCH_3 ; when R^1 and R^2 are both H and R^4 is unsubstituted phenyl, R^3 is not NO_2 ; and when R^1 and R^3 are both H and R^4 is CH_3 , R^2 is not $N(CH_3)_2$.

In embodiments of the invention, compounds of Formula III are those in which R^1 , R^2 and R^3 are each independently selected from H, OH, C_{1-6} alkyl, C_{1-6} alkoxy, NH_2 , $NH-C_{1-6}$ alkyl, $N(C_{1-6}alkyl)(C_{1-6}alkyl)$, SH, $S-C_{1-6}alkyl$, NO_2 , CF_3 , OCF_3 and halo. In preferred embodiments, R^1 , R^2 and R^3 are each independently selected from H, OH, C_{1-4} alkyl, C_{1-4} alkoxy, NH_2 , $NH-C_{1-4}alkyl$, $N(C_{1-4}alkyl)(C_{1-4}alkyl)$, NO_2 , CF_3 , OCF_3 and halo. In more preferred embodiments, R^1 , R^2 and R^3 are each independently selected from H, OH, OCH_3 , NH_2 , $N(CH_3)_2$ and NO_2 . In the most preferred embodiments, R^1 , R^2 and R^3 are each independently selected from H, OH and OCH_3 .

Further embodiments of the invention include compounds of Formula III wherein R^4 is selected from C_{1-6} alkyl, phenyl and pyridyl, wherein phenyl and pyridyl are unsubstituted or substituted with 1-4 substituents, independently selected from C_{1-6} alkyl, C_{1-6} alkoxy and halo. In preferred embodiments of the present invention, R^4 is selected from C_{1-4} alkyl, phenyl and pyridyl, wherein phenyl and pyridyl are unsubstituted or substituted with 1-3 substituents, independently selected from C_{1-4} alkyl, C_{1-4} alkoxy and halo. In more preferred embodiments, R^4 is selected from CH_3 and phenyl, wherein phenyl is unsubstituted or substituted with 1-2 substituents, independently selected from C_{1-4} alkyl, C_{1-4} alkoxy and halo. In the most preferred embodiment R^4 is unsubstituted phenyl.

In a preferred embodiment of the present invention, compounds of Formula I include those in which at least one of R^1 , R^2 and R^3 is OH, more preferably at least two of R^1 , R^2 and R^3 are OH, while R^4 is selected from unsubstituted phenyl and

phenyl substituted with 1-4 substituents, independently selected from C₁₋₆alkyl, C₁₋₆alkoxy and halo.

In specific embodiments of the present invention, the compounds of the invention include:

2-Benzenesulfonyl-5-(3,4-dihydroxyphenyl)-penta-2E,4E-dienitrile (CRVIII-33)

2-Benzenesulfonyl-5-(4-hydroxy-3,5-dimethoxyphenyl)-penta-2E,4E-dienitrile (CRVIII-34)

2-Benzenesulfonyl-5-(4-nitrophenyl)-penta-2E,4E-dienitrile (CRVIII-35)

5-(3,4-Dihydroxyphenyl)-2-(pyridine-2-sulfonyl)-penta-2E,4E-dienitrile (CRVIII-50).

2-(4-Chlorobenzenesulfonyl)-5-(3,4-dihydroxyphenyl)-penta-2E,4E-dienitrile (CRVIII-51)

5-(3,4-Dihydroxyphenyl)-2-(toluene-4-sulfonyl)-penta-2E,4E-dienitrile (CRVIII-52); and

5-(3,4-Dihydroxyphenyl)-2-methanesulfonyl-penta-2E,4E-dienitrile (CRVIII-53).

The present invention includes within its scope prodrugs of the compounds of the invention. In general, such prodrugs will be functional derivatives of a compound of the invention which are readily convertible *in vivo* into the compound from which it is notionally derived. Conventional procedures for the selection and preparation of suitable prodrugs are described, for example, in "Design of Prodrugs" ed. H. Bundgaard, Elsevier, 1985.

The present invention includes radiolabeled forms of compounds of the invention, for example, compounds of the invention labeled by incorporation within the structure ³H or ¹⁴C or a radioactive halogen such as ¹²⁵I.


III. Methods of Preparing Compounds of the Invention

In accordance with another aspect of the present invention, the compounds of the invention can be prepared by processes analogous to those established in the art.

14

Therefore, compounds of this invention may be prepared by the reaction sequence shown in Scheme 1:


Scheme 1



Compounds of the general Formula I or II useful in the practice of this invention can be prepared by Knoevenagel condensation of α,β -unsaturated aldehydes, such as cinnamaldehyde or its various aryl-substituted homologues (V), with an α -cyano ester compound having an active α -methylene group (VI). Similar Knoevenagel condensations using ylidenemalononitriles as active α -methylene group components were described in a review (F. Freeman. Chem. Rev. 1980, V. 80, P. 329-350). Use of a reagent analogous to VI wherein the CO_2X is replaced by SO_2 permits the preparation of compounds of Formulae III and IV. These condensations may, for example, be carried out in a polar solvent, such as ethanol, in the presence of catalytic amounts of a weak base, such as β -alanine. Reaction temperatures may be in the range of 20 to 100 $^\circ\text{C}$, depending on the stability of the materials used in the condensation. Depending on the reactivity of the starting reagents V and/or VI a stronger base, such as piperidine, may also be used to effect the condensation reaction.


Compounds of Formula V may be commercially available, such as cinnamaldehyde, and its 3,5-dimethoxy-4-hydroxy and 4-nitro derivatives. Other compounds of Formula V may be prepared using straightforward procedures. For example, various R^1 , R^2 , R^3 -hydroxy substituted cinnamaldehydes can be prepared from the corresponding commercially available aryl substituted cinnamic acids. Scheme 2 gives an example of the preparation of 3,4-dihydroxycinnamaldehyde (caffeic aldehyde, Va) starting from 3,4-dihydroxycinnamic acid (VII).

15
Scheme 2



Compounds of Formula VI may be prepared, for example, by condensing
5 cyanoacetic acid X with the appropriately substituted alcohol XI under standard conditions, as shown in Scheme 3.

Scheme 3



R¹, R², R³ substituents may be also converted from one functional group to
another, for example by known reduction of nitro groups into amino groups and the
further transformation into dialkylamino groups, or by known conversion of
15 hydroxy groups to halo groups.

In some cases the reactions outlined above may have to be modified, for
instance by use of protective groups, to prevent side reactions due to reactive groups,
such as reactive groups attached as substituents. This may be achieved by means of
20 conventional protecting groups, for example as described in "Protective Groups in
Organic Chemistry" McOmie, J.F.W. Ed., Plenum Press, 1973 and in Greene, T.W.
and Wuts, P.G.M., "Protective Groups in Organic Synthesis", John Wiley & Sons,
1991.

25 The formation of a desired compound salt is achieved using standard
techniques. For example, the neutral compound is treated with an acid or base in a
suitable solvent and the formed salt is isolated by filtration, extraction or any other
suitable method.

The formation of solvates of the compounds of the invention will vary depending on the compound and the solvate. In general, solvates are formed by dissolving the compound in the appropriate solvent and isolating the solvate by cooling or using an antisolvent. The solvate is typically dried or azeotroped under ambient conditions.

Prodrugs of the compounds of the invention may be conventional esters formed with available hydroxy, amino or carboxyl group. For example, when R¹, R² or R³ is OH in a compound as described herein it may be acylated using an activated acid in the presence of a base, and optionally, in inert solvent (e.g., an acid chloride in pyridine). Some common esters which have been utilized as prodrugs are phenyl esters, aliphatic (C₈-C₂₄) esters, acyloxymethyl esters, carbamates and amino acid esters.

A radiolabeled compound of the invention may be prepared using standard methods known in the art. For example, tritium may be incorporated into a compound of the invention using standard techniques, for example by hydrogenation of a suitable precursor to a compound of the invention using tritium gas and a catalyst. Alternatively, a compound of the invention containing radioactive iodo may be prepared from the corresponding trialkyltin (suitably trimethyltin or tributyltin) derivative using standard iodination conditions, such as [¹²⁵I] sodium iodide in the presence of chloramine-T in a suitable solvent, such as dimethylformamide. The trialkyltin compound may be prepared from the corresponding non-radioactive halo, suitably iodo, compound using standard palladium-catalyzed stannylation conditions, for example hexamethylditin in the presence of tetrakis(triphenylphosphine) palladium (0) in an inert solvent, such as dioxane, and at elevated temperatures, suitably 50-100 °C.

IV. Uses

The present invention includes all uses of the compounds as set forth above, including their use in therapeutic methods and compositions for modulating cell

proliferation, preferably inhibiting cell proliferation, their use in diagnostic assays, and their use as research tools.

In one aspect, the present invention provides a method for modulating cell proliferation comprising administering an effective amount of a compound of the invention to a cell or animal in need thereof. Preferably, the invention provides a method of inhibiting cell proliferation comprising administering an effective amount of a compound of the invention to a cell or animal in need thereof. In particular, the method of the invention is useful in inhibiting the proliferation of abnormal but not normal cells. As stated hereinabove, abnormal cells include any type of cell that is causative of or involved in a disease or condition and wherein it is desirable to modulate or inhibit the proliferation of the abnormal cell to treat the disease or condition.

The compounds of the invention may be very effective at killing cancer cells while at the same time not killing normal cells and therefore, may be extremely useful anti-cancer agents. Accordingly, in one embodiment, the present invention provides a method of inhibiting the proliferation of a cancer cell comprising administering an effective amount of a compound of the invention to a cell or animal in need thereof. The present invention also includes a use of a compound or composition of the invention in order to modulate, preferably inhibit, cell proliferation, preferably cancer cell proliferation. The present invention further includes a use of a compound or a composition of the invention to prepare a medicament to modulate, preferably inhibit, cell proliferation, preferably cancer cell proliferation. The cancer cell that can be treated with a compound of the invention may be any type of cancer including, but not limited to, hematopoietic malignancies, such as leukemia.

In a specific embodiment, the present invention provides a method of inhibiting the proliferation of a cell, comprising administering to a cell or animal in need thereof, an effective amount of a compound selected from:

5

2-Cyano-5-(4-hydroxy-3,5-dimethoxyphenyl)-penta-2E,4E-dienoic acid benzyl ester (CRIX-38);

2-Cyano-5-(3,4-dihydroxyphenyl)-penta-2E,4E-dienoic acid benzyl ester (CRIX-39);

10

2-cyano-5-(3,4-dihydroxyphenyl)-penta-2E,4E-dienoic acid 2-[2-(2-methoxyethoxy)ethoxy] ethyl ester (CRIV-42);

2-Cyano-5-(4-hydroxy-3,5-dimethoxyphenyl)-penta-2E,4E-dienoic acid 2-[2-(2-methoxyethoxy)ethoxy]ethyl ester (CRIV-46);

2-Cyano-5-(4-hydroxy-3-methoxyphenyl)-penta-2E,4E-dienoic acid benzyl ester (CRIX-79);

15

2-Benzenesulfonyl-5-(3,4-dihydroxyphenyl)-penta-2E,4E-dienenitrile (CRVIII-33)

2-Benzenesulfonyl-5-(4-hydroxy-3,5-dimethoxyphenyl)-penta-2E,4E-dienenitrile (CRVIII-34)

2-Benzenesulfonyl-5-(4-nitrophenyl)-penta-2E,4E-dienenitrile (CRVIII-35)

20

5-(3,4-Dihydroxyphenyl)-2-(pyridine-2-sulfonyl)-penta-2E,4E-dienenitrile (CRVIII-50)

2-(4-Chlorobenzenesulfonyl)-5-(3,4-dihydroxyphenyl)-penta-2E,4E-dienenitrile (CRVIII-51)

5-(3,4-Dihydroxyphenyl)-2-(toluene-4-sulfonyl)-penta-2E,4E-dienenitrile (CRVIII-

25

52); and

5-(3,4-Dihydroxyphenyl)-2-methanesulfonyl-penta-2E,4E-dienenitrile (CRVIII-53).

The compounds as set forth herein may be effective as *ex vivo* purging agents. For *ex vivo* administration, cells may be removed from a patient and purged *ex vivo* with a compound of the invention. Such a purging should kill the abnormally proliferating cells while leaving normal cells intact. After purging, the treated cells can be washed and reintroduced into the patient.

30

The activity of the compounds of the invention for modulating cell proliferation can be determined using standard assays known to those skilled in the art. Examples of abnormally proliferating cells include malignant or cancerous cells as well as cells such as t-cells that over-proliferate in autoimmune disorders and
5 graft versus host disorders, endothelial and/or epithelial cells that over-proliferate in angiogenesis, and eosinophils that over-proliferate in allergy-related conditions. To assay for efficacy in modulating cell proliferation, the cells may be cultured in the presence of one or more compounds of the invention, and the effect of the compound(s) on the proliferation of the cells can be assayed and compared to the
10 proliferation of the cells in the absence of the compound.

The compounds of the invention may be tyrosine kinase modulators and therefore, useful in modulating tyrosine kinase activity, including the inhibition of tyrosine kinase activity, for the treatment of various conditions such as all
15 proliferative disorders as mentioned above. Accordingly, the invention provides a method of modulating tyrosine kinase activity by administering an effective amount of a compound as set forth herein to a cell or animal in need thereof. In a further aspect, the invention provides a method of inhibiting tyrosine kinase activity by administering an effective amount of a compound as set forth herein to a cell or
20 animal in need thereof.

While the compounds of the invention may act by modulating tyrosine kinase activity, one of skill in the art will appreciate that other modes or mechanisms of action for the compounds of the invention are possible.
25

The compounds as set forth herein are preferably formulated into pharmaceutical compositions for administration to human subjects in a biologically compatible form suitable for administration *in vivo*. Accordingly, in another aspect, the present invention provides a pharmaceutical composition comprising a
30 compound of the invention in admixture with a suitable diluent or carrier.

The compositions containing the compounds of the invention can be prepared by known methods for the preparation of pharmaceutically acceptable compositions which can be administered to subjects, such that an effective quantity of the active substance is combined in a mixture with a pharmaceutically acceptable vehicle.

5 Suitable vehicles are described, for example, in Remington's Pharmaceutical Sciences (Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa., USA 1985). On this basis, the compositions include, albeit not exclusively, solutions of the substances in association with one or more pharmaceutically acceptable vehicles or diluents, and contained in buffered
10 solutions with a suitable pH and iso-osmotic with the physiological fluids.

The compounds as set forth herein may be used in the form of the free base, in the form of salts, solvates and as hydrates. All forms are within the scope of the invention. Acid addition salts may be formed and provide a more convenient form
15 for use; in practice, use of the salt form inherently amounts to use of the base form. The acids which can be used to prepare the acid addition salts include preferably those which produce, when combined with the free base, pharmaceutically acceptable salts, that is, salts whose anions are non-toxic to the animal organism in pharmaceutical doses of the salts, so that the beneficial properties inherent in the
20 free base are not vitiated by side effects ascribable to the anions. Although pharmaceutically acceptable salts of the basic compounds are preferred, all acid addition salts are useful as sources of the free base form even if the particular salt per se is desired only as an intermediate product as, for example, when the salt is formed only for the purposes of purification and identification, or when it is used as
25 an intermediate in preparing a pharmaceutically acceptable salt by ion exchange procedures.

Pharmaceutically acceptable salts within the scope of the invention include those derived from the following acids; mineral acids such as hydrochloric acid,
30 sulfuric acid, phosphoric acid and sulfamic acid; and organic acids such as acetic acid, citric acid, lactic acid, tartaric acid, malonic acid, methanesulfonic acid,

21

ethanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, cyclohexylsulfamic acid, quinic acid, and the like.

In accordance with the methods of the invention, the described compounds or salts or solvates thereof may be administered to a patient in a variety of forms depending on the selected route of administration, as will be understood by those skilled in the art. The compositions of the invention may be administered orally or parenterally. Parenteral administration includes intravenous, intraperitoneal, subcutaneous, intramuscular, transepithelial, nasal, intrapulmonary, intrathecal, rectal and topical modes of administration. Parenteral administration may be by continuous infusion over a selected period of time.

A compound of the invention or a salt or solvate thereof may be orally administered, for example, with an inert diluent or with an assimilable edible carrier, or it may be enclosed in hard or soft shell gelatin capsules, or it may be compressed into tablets, or it may be incorporated directly with the food of the diet. For oral therapeutic administration, the compound of the invention may be incorporated with excipient and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like.

20

A compound of the invention may also be administered parenterally or intraperitoneally. Solutions of a compound of the invention as a free base or pharmacologically acceptable salt or solvate can be prepared in water suitably mixed with a surfactant such as hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, DMSO and mixtures thereof with or without alcohol, and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms. A person skilled in the art would know how to prepare suitable formulations. Conventional procedures and ingredients for the selection and preparation of suitable formulations are described, for example, in Remington's Pharmaceutical Sciences (1990 - 18th edition) and in The United States Pharmacopeia: The National Formulary (USP 24 NF19) published in 1999.

22

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersion and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases the form must be sterile and must be fluid to the extent that easy syringability exists.

5

The compounds as set forth herein may be administered to an animal alone or in combination with pharmaceutically acceptable carriers, as noted above, the proportion of which is determined by the solubility and chemical nature of the compound, chosen route of administration and standard pharmaceutical practice.

10

The dosage of the compounds and/or compositions as set forth herein can vary depending on many factors such as the pharmacodynamic properties of the compound, the mode of administration, the age, health and weight of the recipient, the nature and extent of the symptoms, the frequency of the treatment and the type of concurrent treatment, if any, and the clearance rate of the compound in the animal to be treated. One of skill in the art can determine the appropriate dosage based on the above factors. The compounds as set forth herein may be administered initially in a suitable dosage that may be adjusted as required, depending on the clinical response.

20

The compounds as set forth herein may be used alone or in combination with other agents that modulate tyrosine kinase activity or in combination with other types of treatment (which may or may not modulate tyrosine kinase activity) for cell proliferative disorders. Agents known in the art that inhibit tyrosine kinase activity include, but are not limited to, antisense nucleic acid and ribozymes targeted to nucleic acid encoding a receptor tyrosine kinase, antibodies able to modulate tyrosine kinase activity and other small molecule tyrosine kinase inhibitors such as those described in US 5,891,917, US 5,217,999, US 5,773,476, US 5,935,993, US 5,656,655, US 5,677,329 and US 5,789,427.

30

In addition to the above-mentioned therapeutic uses, the compounds as set forth herein are also useful in diagnostic assays, screening assays and as research tools.

5 In diagnostic assays the compounds as set forth herein may be useful in identifying or detecting a cell proliferative disorder. In such an embodiment, the compounds as set forth herein may be radiolabelled (as hereinbefore described) and contacted with a population of cells. The presence of the radiolabelled compound on the cells may indicate a cell proliferative disorder.

10

In screening assays, the compounds as set forth herein may be used to identify other compounds that modulate cell proliferation or tyrosine kinase activity. As research tools, the compounds of the invention may be used in receptor binding assays and assays to study the localization of tyrosine kinases. In such assays, the
15 compounds may also be radiolabelled.

The following non-limiting examples are illustrative of the present invention:

Examples

20 *Materials and Methods for Examples 1-5*

¹H NMR spectra were obtained on a Varian Unity Plus spectrometer at 500 MHz in acetone-d₆ with Me₄Si as an internal standard ($\delta=0$). Electrospray mass spectra were acquired on an API III+ triple quadrupole mass spectrometer (PE Sciex, Thornhill, Canada). Samples were directly introduced into the electrospray
25 ionization source using an HPLC pump. Thin layer chromatography was performed with UV-254 aluminum-backed TLC sheets of 0.25 mm thickness (Kieselgel 60 F254, Merck, Germany). HPLC chromatograms and UV spectra were obtained on a model 600 liquid chromatograph (Waters, USA) with a model 996 PDA detector. UV spectra of the compounds were recorded from 200 to 800 nm in steps of 1.2 nm.
30 Spectra were obtained in the mobile phase in which they were eluted from the column. The column was 5 μ m Zorbax SB-CN (Agilent Technologies, USA). Gradient separations were conducted using the following buffers: (A) 0.1% H₃PO₄

24

in H₂O; (B) MeCN-H₂O, 4:1 + 0.1% H₃PO₄. The following gradient system was employed: 20% of the buffer B in the buffer A for 10 min, followed by linearly increasing the percentage of the buffer B from 20 to 90% during 20 min. The flow rate was 1.0 mL/min. Vacuum distillations were done using Kugelrohr apparatus
5 (Aldrich) at stated temperatures of an oven.

4-Hydroxy-3,5-dimethoxycinnamaldehyde and 4-hydroxy-3-methoxycinnamaldehyde, as well as all other reagents were purchased from Aldrich (USA) and were used as received. Solvents were purchased from Caledon
10 (Canada). A modified synthetic scheme [see C.M. Roifman, T. Grunberger, O. Rounova, P. Demin, and N. Sharfe. PCT Int. Appl. WO 2001079158, issued 10/25/2001, 100 pp.] was used for the preparation of caffeoyl aldehyde. The direct precursor of the aldehyde, 3,4-bis-(*t*-butyldimethylsilyloxy)cinnamyl alcohol, was obtained as described in an article [M. Potgieter, G.L. Wenteler, and S.E. Drewes.
15 Phytochemistry, 1988, V. 27, No. 4, P. 1101-1104].

Example 1: 3,4-Dihydroxycinnamaldehyde (caffeoyl aldehyde)

3,4-Bis-(*t*-butyldimethylsilyloxy)cinnamyl alcohol (7.88 g, 20 mmol) was dissolved in 1000 mL CH₂Cl₂, 17.2 g activated MnO₂ (200 mmol) were added and
20 the mixture was vigorously stirred for 24 h at 20 °C. MnO₂ was filtered off and the filtrate was taken to dryness. To the obtained 3,4-bis(tBDMS)caffeoyl aldehyde 250 mL of CHCl₃ was added followed by addition of *n*-Bu₄NF monohydrate (11.6 g, 40 mmol). The mixture was stirred at 20 °C for 30 min and worked up with 300 mL of 5% HCl. Chloroform layer was separated, washed with H₂O, dried with Na₂SO₄ and
25 taken to dryness. The residue was purified on a silica gel column, eluent MeOH-CHCl₃, 1:4 + 1% AcOH. The solvents were evaporated and the crystalline residue was washed with CHCl₃ to give caffeoyl aldehyde (1.77 g, 54%). M.p. 200-202 °C, λ_{max} 222, 239, 249, 342 nm. ¹H-NMR (δ, ppm): 6.54 (dd, 1H, J 7.7 and 15.8 Hz, H_α), 6.91 (d, 1H, J 8.2 Hz, H₅), 7.12 (br.d, 1H, J 8.2 Hz, H₆), 7.21 (br.s, 1H, H₂),
30 7.52 (d, 1H, J 15.8 Hz, H_β), 9.62 (d, 1H, J 7.7 Hz, CHO).

Example 2: Cyanoacetic acid benzyl ester (IVa)

To a mixture of cyanoacetic acid (VII) 2.34 g (27.5 mmol) and benzyl alcohol (VIIIa) 2.70 g (25 mmol) in 60 mL of toluene 50 mg of *p*-TsOH was added and the mixture was refluxed in a flask with a Dean-Stark trap for 24 h. The mixture was cooled down and toluene was washed with H₂O and toluene was evaporated. The residue was distilled *in vacuo* (Kugelrohr apparatus, 0.1 mm Hg, T. oven 170-180 °C). Yield 3.36 g (77%), a colorless oil. MS (*m/z*, rel. intensity, %): 167.0 ([M+NH₄-CN]⁺, 27), 193.0 ([M+NH₄]⁺, 100).

10

Example 3: Cyanoacetic acid 2-[2-(2-methoxyethoxy)ethoxy] ester (IVb)

To a mixture of cyanoacetic acid (VII) (9.4 g, 110 mmol) and 2-[2-(2-methoxyethoxy)ethoxy]ethanol (VIIIb) (16.4 g, 100 mmol) in 60 mL of toluene 50 mg of *p*-TsOH was added and the mixture was refluxed in a flask with a Dean-Stark trap for 24 h. The mixture was cooled down and toluene was washed with H₂O and toluene was evaporated. The compound was distilled *in vacuo* (Kugelrohr apparatus, 0.1 mm Hg, T. oven 210 °C). Yield 19.25 g (83%), a colorless oil. MS (*m/z*, rel. intensity, %): 232.0 ([M+H]⁺, 100), 249.2 ([M+NH₄]⁺, 87).

Example 4: Knoevenagel condensation of substituted acetonitriles with hydroxyl-substituted cinnamaldehydes

To 0.2 mmol of 3,4-dihydroxycinnamaldehyde in 3 mL EtOH a corresponding substituted acetonitrile or the like was added followed by addition of 0.4 mmol of piperidine. A dark red solution was stirred at 20 °C for 0.5 h. 0.5 mL 1 N HCl and 5 mL of H₂O were added and EtOH was evaporated with the stream of N₂. The precipitated powder was filtered off, washed with H₂O and dried in a desiccator *in vacuo* in the presence of solid NaOH. Yields were 55-70%. For a list of obtained products (CRIX-38, CRIX-39, CRIV-42, CRIV-46, CRIX-79, CRVIII-33, CRVIII-34, CRVIII-50, CRVIII-51, CRVIII-52, CRVIII-53) see Tables 1 and 5. For the mass spectra and UV maxima see Tables 2 and 6. For the NMR data see Tables 3 and 7.

30

Example 5: Knoevenagel condensation of substituted acetonitriles with nitro-substituted cinnamaldehydes

To 0.2 mmol of 4-nitrocinnamaldehyde in 3 mL EtOH a corresponding substituted acetonitrile was added followed by addition of 0.5-1.0 mg of β -alanine.

5 A deep yellow solution was stirred at 90 °C for 4 h. 0.5 mL 1 N HCl and 5 mL of H₂O were added and EtOH was evaporated with the stream of N₂. The precipitated powder was filtered off, and washed with H₂O. The obtained yellow powder was recrystallized from EtOH-H₂O, the crystals were washed with H₂O and dried in a desiccator *in vacuo* in the presence of solid NaOH. Yields were 50-60%. For the list

10 of obtained products (CRVIII-35) see Table 5. For the mass spectra and UV maxims see Table 6. For the NMR data see Table 7.

Example 6: Effect of Test Compounds Upon Normal Bone Marrow Differentiation in Culture

15 A CFU-GEMM assay may be performed according to Fauser and Messner (1978, Blood, 52(6) 143-8) and Messner and Fauser (1980, Blut, 41(5) 327-33).

Example 7: Killing of Philadelphia Positive Acute Lymphoblastic Leukemia by Low-dose Test Compounds in Culture.

20 Ph+ ALL cells are plated in 1 ml volumes, in the absence of exogenous growth factors, into 35 mm petri dishes (Nunc, Gibco) containing alpha MEM (Gibco) plus 10% FCS (Cansera Rexdale, ON.) in 0.9% (vol/vol) methylcellulose (Fluka, Switzerland) with various concentrations of test compounds. Cultures are set at 37 °C, 5% CO₂ in a humidified atmosphere. Colonies consisting of more than 20

25 cells are counted at 12 days or earlier using an inverted microscope.

Example 8: Killing of Philadelphia positive Z119 Acute Lymphoblastic Leukemia Cells by Low-dose Test Compounds in Culture.

Z119 cells are plated in 1 ml volumes at a density of 1×10^4 cells/ml, in the

30 absence of exogenous growth factors, into 35 mm petri dishes (Nunc, Gibco) containing IMDM (OCI, Toronto) plus 20% FCS (Cansera Rexdale, ON) in 0.9% (vol/vol) methylcellulose (Fluka, Switzerland) with various concentrations of test

compounds. Cultures are set at 37 °C, 5% CO₂ in a humidified atmosphere. Colonies consisting of more than 20 cells are counted at 7 days or earlier using an inverted microscope. Results see Tables 4 and 8.






5 Example 9: Killing of AML-3 Acute Myeloid Leukemia Cells by Low-dose Test Compounds in Culture

OCI-AML-3 cells are plated in 35mm petri dishes (Nunc, Gibco) in 1 ml volumes at a density of 3.3×10^3 cells/ml, in the absence of exogenous growth factors, containing alpha MEM plus 20% FCS (Cansera, Rexdale Ont.), and 0.9%
10 (vol/vol) methylcellulose (Fluka, Switzerland) and various concentrations of test compounds. Cell cultures are incubated in a humidified atmosphere at 37 °C with 5% CO₂. Colonies containing more than 20 cells are scored, using an inverted microscope, at 5-6 days. Results see Tables 4 and 8.

15 While the present invention has been described with reference to what are presently considered to be the preferred examples, it is to be understood that the invention is not limited to the disclosed examples. To the contrary, the invention is intended to cover various modifications and equivalent arrangements included within the spirit and scope of the appended claims.

20 All publications, patents and patent applications are herein incorporated by reference in their entirety to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference in its entirety.

28
Table 1

Formula	Chemical name	Code
	2-Cyano-5-(4-hydroxy-3,5-dimethoxyphenyl)-penta-2E,4E-dienoic acid benzyl ester	CRIX-38
	2-Cyano-5-(3,4-dihydroxyphenyl)-penta-2E,4E-dienoic acid benzyl ester	CRIX-39
	2-Cyano-5-(3,4-dihydroxyphenyl)-penta-2E,4E-dienoic acid 2-[2-(2-methoxyethoxy)ethoxy]ethyl ester	CRIV-42
	2-Cyano-5-(4-hydroxy-3,5-dimethoxyphenyl)-penta-2E,4E-dienoic acid 2-[2-(2-methoxyethoxy)ethoxy]ethyl ester	CRIV-46
	2-Cyano-5-(4-hydroxy-3-methoxyphenyl)-penta-2E,4E-dienoic acid benzyl ester	CRIX-79

29
Table 2

Code	MS, m/z (rel.intensity, %)	λ_{max} , nm
CRIX-38	366.0 ($[\text{M}+\text{H}]^+$, 82), 383.0 ($[\text{M}+\text{NH}_4]^+$, 100)	267, 407
CRIX-39	322.2 ($[\text{M}+\text{H}]^+$, 59), 339.0 ($[\text{M}+\text{NH}_4]^+$, 100)	265, 405
CRIV-42	378.0 ($[\text{M}+\text{H}]^+$, 28), 395.1 ($[\text{M}+\text{NH}_4]^+$, 100)	266, 401
CRIV-46	422.2 ($[\text{M}+\text{H}]^+$, 71), 439.2 ($[\text{M}+\text{NH}_4]^+$, 100)	267, 406
CRIX-79	335.8 ($[\text{M}+\text{H}]^+$, 100), 353.0 ($[\text{M}+\text{NH}_4]^+$, 99)	263, 400








30
Table 3

Code	H ²	H ³	H ⁴	H ⁵	H ⁶	PhCH= C	PhC=C H	CH=CC N	other
CRIX-38	7.08 (br.s, 1H)	-	-	-	7.08 (br.s, 1H)	7.52 (d, 1H, J 15.1 Hz)	7.18 (dd, 1H, J 11.8 and 15.1 Hz)	8.08 (d, 1H, J 11.8 Hz)	5.30 (s, 2H, COOCH ₂ P h', 7.32- 7.46 (m, 5H, Ph')
CRIX-39	7.22 (d, 1H, J 2.1 Hz)	-	-	6.88 (d, 1H, J 8.1 Hz)	7.09 (dd, 1H, J 2.1 and 8.1 Hz)	7.47 (d, 1H, J 15.1 Hz)	7.06 (dd, 1H, J 11.5 and 15.1 Hz)	8.08 (d, 1H, J 11.5 Hz)	5.30 (s, 2H, COOCH ₂ P h', 7.32- 7.45 (m, 5H, Ph')
CRIV-42	7.25 (br.s, 1H)	-	-	6.90 (d, 1H, J 8.0 Hz)	7.12 (br.d, 1H, J 8.0 Hz)	7.49 (d, 1H, J 15.1 Hz)	7.07 (dd, 1H, J 11.8 and 15.1 Hz)	8.11 (d, 1H, J 11.8 Hz)	3.29 (s, 3H, Me), 3.50; 3.60; 3.66; 3.79, 4.40 (5 × m, 12H, (CH ₂ CH ₂ O) ₂)
CRIV-46	7.12 (br.s, 1H)	-	-	-	7.12 (br.s, 1H)	7.52 (d, 1H, J 15.3 Hz)	7.21 (dd, 1H, J 11.7 and 15.3 Hz)	8.06 (d, 1H, J 11.7 Hz)	3.28 (s, 3H, Me), 3.90 (s, 6H, 3- and 5-OMe), 3.47; 3.58; 3.65; 3.77; 4.38 (5 × m, 12H, (CH ₂ CH ₂ O) ₂)
CRIX-79	7.43 (d, 1H, J 2.0 Hz)	-	-	6.92 (d, 1H, J 8.3 Hz)	7.24 (dd, 1H, J 2.0 and 8.3 Hz)	7.54 (d, 1H, J 15.3 Hz)	7.19 (dd, 1H, J 11.7 and 15.3 Hz)	8.10 (d, 1H, J 11.7 Hz)	3.93 (s, 3H, 3- OMe), 5.32 (s, 2H, COOCH ₂ P h'), 7.36- 7.43; 7.47 (2 × m, 5H, Ph')

31
Table 4

Compound	ALL (Z 119) IC 50, μ M	AML (OCI AML-3) IC 50, μ M	NBM (BFU-E) IC 50, μ M
CRIX-38	> 1.0	> 0.5	\leq 10.0
CRIX-39	\geq 0.6	= 0.09	= 8.0
CRIV-42	\geq 0.5	= 0.09	\geq 10.0
CRIV-46	> 0.5	\geq 0.5	\geq 10.0

32
Table 5

Formula	Chemical name	Code
	2-Benzenesulfonyl-5-(3,4-dihydroxyphenyl)-penta-2E,4E-dienitrile	CRVIII-33
	2-Benzenesulfonyl-5-(4-hydroxy-3,5-dimethoxyphenyl)-penta-2E,4E-dienitrile	CRVIII-34
	2-Benzenesulfonyl-5-(4-nitrophenyl)-penta-2E,4E-dienitrile	CRVIII-35
	5-(3,4-Dihydroxyphenyl)-2-(pyridine-2-sulfonyl)-penta-2E,4E-dienitrile	CRVIII-50
	2-(4-Chlorobenzenesulfonyl)-5-(3,4-dihydroxyphenyl)-penta-2E,4E-dienitrile	CRVIII-51
	5-(3,4-Dihydroxyphenyl)-2-(toluene-4-sulfonyl)-penta-2E,4E-dienitrile	CRVIII-52
	5-(3,4-Dihydroxyphenyl)-2-methanesulfonyl-penta-2E,4E-dienitrile	CRVIII-53

33
Table 6

Code	MS, m/z (rel.intensity, %)	λ_{\max} , nm
CRVIII-33	242.4 (21), 328.0 ($[M+H]^+$, 100), 345.0 ($[M+NH_4]^+$, 40)	266, 406
CRVIII-34	372.0 ($[M+H]^+$, 100), 389.0 ($[M+NH_4]^+$, 64)	267, 413
CRVIII-35	341.0 ($[M+H]^+$, 35), 358.0 ($[M+NH_4]^+$, 100)	350
CRVIII-50	329.0 ($[M+H]^+$, 100), 346.0 ($[M+NH_4]^+$, 12)	265, 415
CRVIII-51	361.8 ($[M+H]^+$, 100), 378.8 ($[M+NH_4]^+$, 87)	265, 412
CRVIII-52	341.8 ($[M+H]^+$, 19), 359.0 ($[M+NH_4]^+$, 100)	265, 406
CRVIII-53	265.8 ($[M+H]^+$, 5), 282.8 ($[M+NH_4]^+$, 100)	262, 399

34
Table 7


Code	H ²	H ³	H ⁴	H ⁵	H ⁶	PhCH= C	PhC=C H	CH=CC N	other
CRVIII-33	7.22 (d, 1H, J 2.2 Hz)	-	-	6.88 (d, 1H, J 8.3 Hz)	7.11 (dd, 1H, J 2.2 and 8.3 Hz)	7.58 (d, 1H, J 15.0 Hz)	6.93 (dd, 1H, J 11.7 and 15.0 Hz)	8.10 (d, 1H, J 11.7 Hz)	7.70; 7.78; 7.96 (3 × m, 5H, Ph')
CRVIII-34	7.14 (br.s, 1H)	-	-	-	7.14 (br.s, 1H)	7.66 (d, 1H, J 15.0 Hz)	7.11 (dd, 1H, J 11.6 and 15.0 Hz)	8.12 (d, 1H, J 11.6 Hz)	7.74; 7.82; 7.99 (3 × m, 5H, Ph')
CRVIII-35	8.08 (m, 2H, + H6)	8.32 (m, 2H, + H5)	-	8.32 (m, 2H, + H3)	8.08 (m, 2H, + H2)	7.88 (d, 1H, J 15.3 Hz)	7.42 (dd, 1H, J 11.4 and 15.3 Hz)	8.28 (d, 1H, J 11.4 Hz)	7.73; 7.86; 8.04 (3 × m, 5H, Ph')
CRVIII-50	7.30 (d, 1H, J 2.1 Hz)	-	-	6.94 (d, 1H, J 8.3 Hz)	7.20 (dd, 1H, J 2.1 and 8.3 Hz)	7.68 (d, 1H, J 15.1 Hz)	7.05 (dd, 1H, J 11.6 and 15.1 Hz)	8.14 (d, 1H, J 11.6 Hz)	7.78; 8.17-8.24; 8.79 (3 × m, 4H, Py)
CRVIII-51	7.28 (d, 1H, J 2.2 Hz)	-	-	6.93 (d, 1H, J 8.3 Hz)	7.18 (dd, 1H, J 2.2 and 8.3 Hz)	7.63 (d, 1H, J 15.1 Hz)	6.98 (dd, 1H, J 11.6 and 15.1 Hz)	8.14 (d, 1H, J 11.6 Hz)	7.76; 8.00 (2 × m, 4H, Ph')
CRVIII-52	7.26 (d, 1H, J 2.1 Hz)	-	-	6.92 (d, 1H, J 8.3 Hz)	7.16 (dd, 1H, J 2.1 and 8.3 Hz)	7.60 (d, 1H, J 15.1 Hz)	6.96 (dd, 1H, J 11.6 and 15.1 Hz)	8.09 (d, 1H, J 11.6 Hz)	3.58 (s, 3H, Ph'CH ₃), 7.52; 7.86 (2 × m, 4H, Ph')
CRVIII-53	7.29 (d, 1H, J 2.2 Hz)	-	-	6.93 (d, 1H, J 8.2 Hz)	7.17 (dd, 1H, J 2.2 and 8.3 Hz)	7.57 (d, 1H, J 15.0 Hz)	7.01 (dd, 1H, J 11.5 and 15.0 Hz)	7.94 (d, 1H, J 11.5 Hz)	3.23 (s, 3H, SO ₂ Me)

35
Table 8

Compound	ALL (Z 119) IC 50, μM	AML (OCI AML-3) IC 50, μM	NBM (BFU-E) IC 50, μM
CRVIII-33	≥ 0.55	> 0.5	$= 12.0$
CRVIII-34	> 1.0	> 0.5	≤ 10.0
CRVIII-35	> 1.0	> 0.5	> 5.0
CRVIII-50	≥ 0.17	≥ 0.25	≤ 5.0
CRVIII-51	$= 0.23$	0.25-0.5	≥ 5.0
CRVIII-52	0.25-0.5	0.25-0.5	≤ 10.0
CRVIII-53	0.45	0.25-0.5	≤ 10.0

Claims:

1. A compound of Formula I, or a salt, solvate or hydrate thereof:



5 wherein

R^1 , R^2 and R^3 are each independently selected from H, OH, C_{1-6} alkyl, C_{1-6} alkoxy, NH_2 , $NH-C_{1-6}$ alkyl, $N(C_{1-6}$ alkyl)(C_{1-6} alkyl), SH, $S-C_{1-6}$ alkyl, NO_2 , CF_3 , OCF_3 and halo;

R^4 is unsubstituted Ar, or Ar substituted with 1-4 substituents, independently selected
10 from C_{1-6} alkyl, C_{1-6} alkoxy and halo;

X is selected from $(CH_2CH_2O)_n$ and $(CH_2)_n$, and

$n = 1-4$.

2. The compound according to claim 1, wherein

R^1 , R^2 and R^3 are each independently selected from H, OH, C_{1-4} alkyl, C_{1-4} alkoxy, NH_2 , $NH-C_{1-4}$ alkyl, $N(C_{1-4}$ alkyl)(C_{1-4} alkyl), NO_2 , CF_3 , OCF_3 and halo;

15 R^4 is C_{1-6} alkyl,

R^4 is C_{1-6} alkyl,

X is $(CH_2CH_2O)_n$, and

$n = 1-4$.


3. The compound according to any of claims 1 and 2, wherein R^1 , R^2 and R^3 are
20 each independently selected from H, OH, C_{1-4} alkyl, C_{1-4} alkoxy, NH_2 , $NH-C_{1-4}$ alkyl, $N(C_{1-4}$ alkyl)(C_{1-4} alkyl), NO_2 , CF_3 , OCF_3 and halo.

4. The compound according to claim 3, wherein R^1 , R^2 and R^3 are each
independently selected from H, OH, OCH_3 , NH_2 , $N(CH_3)_2$ and NO_2 .

5. The compound according to claim 4, wherein R^1 , R^2 and R^3 are each
25 independently selected from H, OH, and OCH_3 .

6. The compound according to claim 1, wherein R^4 is unsubstituted Ar.
7. The compound according to claim 6, wherein R^4 is phenyl.
8. The compound according to claim 2, wherein R^4 is methyl or ethyl.
9. The compound according to claim 8, wherein R^4 is methyl.
- 5 10. The compound according to claim 9, wherein n is 2-3.
11. The compound according to claim 10, wherein n is 3.
12. A compound selected from:
2-Cyano-5-(4-hydroxy-3,5-dimethoxyphenyl)-penta-2E,4E-dienoic acid benzyl ester
(CRIX-38)
10 2-Cyano-5-(3,4-dihydroxyphenyl)-penta-2E,4E-dienoic acid benzyl ester (CRIX-39)
2-Cyano-5-(3,4-dihydroxyphenyl)-penta-2E,4E-dienoic acid 2-[2-(2-methoxyethoxy)ethoxy] ethyl ester (CRIV-42)
2-Cyano-5-(4-hydroxy-3,5-dimethoxyphenyl)-penta-2E,4E-dienoic acid 2-[2-(2-methoxyethoxy)ethoxy]ethyl ester (CRIV-46); and
15 2-Cyano-5-(4-hydroxy-3-methoxyphenyl)-penta-2E,4E-dienoic acid benzyl ester
(CRIX-79).
13. A composition comprising a compound according to any one of claims 1 to 12 in admixture with a pharmaceutically acceptable diluent or carrier.
14. A use of a compound according to any of claims 1-12, and/or a composition
20 according to claim 13, to prepare a medicament to modulate cell proliferation.
15. The use according to claim 14, for inhibiting cell proliferation.
16. The use according to claim 15, wherein the cell is a malignant hematopoietic cell.

17. A method of modulating cell proliferation comprising administering an effective amount of a compound according to any of claims 1-12, and/or a composition according to claim 13, to a cell or animal in need thereof.
18. The method according to claim 17, for inhibiting cell proliferation.
19. The method according to claim 18 wherein the cell is a malignant hematopoietic cell.
20. A compound of Formula III, and/or a salt, solvate, or hydrate thereof:




wherein

- 10 R^1 , R^2 and R^3 are each independently selected from H, OH, C_{1-6} alkyl, C_{1-6} alkoxy, NH_2 , $NH-C_{1-6}$ alkyl, $N(C_{1-6}$ alkyl)(C_{1-6} alkyl), SH, $S-C_{1-6}$ alkyl, NO_2 , CF_3 , OCF_3 and halo; and R^4 is selected from C_{1-6} alkyl, phenyl and pyridyl, wherein phenyl and pyridyl are unsubstituted or substituted with 1-4 substituents, independently selected from C_{1-6} alkyl, C_{1-6} alkoxy and halo, with the provisos that when R^1 and R^3 are both H and R^4 is unsubstituted phenyl, R^2 is not H, Cl, or OCH_3 ; when R^1 and R^2 are both H and R^4 is unsubstituted phenyl, R^3 is not NO_2 ; and when R^1 and R^3 are both H and R^4 is CH_3 , R^2 is not $N(CH_3)_2$.
- 15
21. The compound according to claim 1, wherein R^1 , R^2 and R^3 are each independently selected from H, OH, C_{1-4} alkyl, C_{1-4} alkoxy, NH_2 , $NH-C_{1-4}$ alkyl, $N(C_{1-4}$ alkyl)(C_{1-4} alkyl), NO_2 , CF_3 , OCF_3 and halo.
- 20
22. The compound according to claim 21, R^1 , R^2 and R^3 are each independently selected from H, OH, OCH_3 , NH_2 , $N(CH_3)_2$ and NO_2 .

23. The compound according to claim 20, wherein R⁴ is selected from C₁₋₄alkyl, phenyl and pyridyl.
24. The compound according to claim 23, wherein R⁴ is selected from CH₃ and phenyl.
- 5 25. The compound according to claim 24, wherein R⁴ is unsubstituted phenyl.
26. The compound according to claim 20, wherein phenyl and pyridyl are unsubstituted or substituted with 1-3 substituents, independently selected from C₁₋₄alkyl, C₁₋₄alkoxy and halo.
27. The compound according to claim 24, wherein phenyl is unsubstituted or
10 substituted with 1-2 substituents, independently selected from C₁₋₄alkyl, C₁₋₄alkoxy and halo.
28. The compound according to claim 20, wherein at least one of R¹, R² and R³ is OH while R⁴ is selected from unsubstituted phenyl and phenyl substituted with 1-4 substituents, independently selected from C₁₋₄alkyl, C₁₋₄alkoxy and halo.
- 15 29. A compound selected from:
2-Benzenesulfonyl-5-(3,4-dihydroxyphenyl)-penta-2E,4E-dienenitrile (CRVIII-33),
2-Benzenesulfonyl-5-(4-hydroxy-3,5-dimethoxyphenyl)-penta-2E,4E-dienenitrile
(CRVIII-34),
2-Benzenesulfonyl-5-(4-nitrophenyl)-penta-2E,4E-dienenitrile (CRVIII-35),
20 5-(3,4-Dihydroxyphenyl)-2-(pyridine-2-sulfonyl)-penta-2E,4E-dienenitrile (CRVIII-50),
2-(4-Chlorobenzenesulfonyl)-5-(3,4-dihydroxyphenyl)-penta-2E,4E-dienenitrile
(CRVIII-51),
5-(3,4-Dihydroxyphenyl)-2-(toluene-4-sulfonyl)-penta-2E,4E-dienenitrile (CRVIII-52), and
25 5-(3,4-Dihydroxyphenyl)-2-methanesulfonyl-penta-2E,4E-dienenitrile (CRVIII-53).

30. A composition comprising a compound according to any one of claims 20 to 29 in admixture with a pharmaceutically acceptable diluent or carrier.

31. A composition comprising, in admixture with a pharmaceutically acceptable diluent or carrier, a compound of Formula IV, and/or a salt, solvate or hydrate thereof:




wherein

R^1 , R^2 and R^3 are each independently selected from H, OH, C_{1-6} alkyl, C_{1-6} alkoxy, NH_2 , $NH-C_{1-6}$ alkyl, $N(C_{1-6}$ alkyl)(C_{1-6} alkyl), SH, $S-C_{1-6}$ alkyl, NO_2 , CF_3 , OCF_3 and halo; and

R^4 is selected from C_{1-6} alkyl, phenyl and pyridyl, wherein phenyl and pyridyl are unsubstituted or substituted with 1-4 substituents, independently selected from C_{1-6} alkyl, C_{1-6} alkoxy and halo.

32. A use to prepare a medicament to modulate cell proliferation of a composition according to claim 30 or 31, and/or a compound capable of modulating cell proliferation of Formula IV, and/or a salt, solvate or hydrate thereof:




wherein

R^1 , R^2 and R^3 are each independently selected from H, OH, C_{1-6} alkyl, C_{1-6} alkoxy, NH_2 , $NH-C_{1-6}$ alkyl, $N(C_{1-6}$ alkyl)(C_{1-6} alkyl), SH,

$S-C_{1-6}$ alkyl, NO_2 , CF_3 , OCF_3 and halo; and

R^4 is selected from C_{1-6} alkyl, phenyl and pyridyl, wherein phenyl and pyridyl are unsubstituted or substituted with 1-4 substituents, independently selected from C_{1-6} alkyl, C_{1-6} alkoxy and halo.

33. The use according to claim 13, for inhibiting cell proliferation.
34. The use according to claim 14 wherein the cell is a malignant hematopoietic cell.
35. A method of modulating cell proliferation comprising administering to a cell
5 or animal in need thereof an effective amount of a composition according to any of claims 30 and 31, and/or a compound capable of modulating cell proliferation of Formula IV, and/or a salt, solvate or hydrate thereof:



wherein

- 10 R¹, R² and R³ are each independently selected from H, OH, C₁₋₆alkyl, C₁₋₆alkoxy, NH₂, NH-C₁₋₆alkyl, N(C₁₋₆alkyl)(C₁₋₆alkyl), SH, S-C₁₋₆alkyl, NO₂, CF₃, OCF₃ and halo; and
- R⁴ is selected from C₁₋₆alkyl, phenyl and pyridyl, wherein phenyl and pyridyl are unsubstituted or substituted with 1-4 substituents, independently selected from
15 C₁₋₆alkyl, C₁₋₆alkoxy and halo.
36. The method according to claim 35, for inhibiting cell proliferation.
37. The method according to claim 36, wherein the cell is a malignant hematopoietic cell.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 03/00032

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C255/41 C07C309/73 A61K31/165

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 01 79158 A (GRUNBERGER THOMAS ; DEMIN PETER (CA); ROIFMAN CHAIM M (CA); ROUNOVA) 25 October 2001 (2001-10-25) cited in the application the whole document	1-19
A	US 5 789 427 A (GAZIT AVIV ET AL) 4 August 1998 (1998-08-04) cited in the application	20-29
Y	column 5, line 4 - line 24 column 43, line 18 - line 30 table I, col.11; table VI, col. 39; fig.2b; fig.2d; fig.3g, M26;fig.3h, M30-M31; fig.3i, M32-M34 column 43, line 55 -column 44, line 13	25-37
	--- -/-- ---	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

15 Apr11 2003

Date of mailing of the international search report

13/05/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Seelmann, M

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/CA 03/00032

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4 632 895 A (PATEL RANJANA C ET AL) 30 December 1986 (1986-12-30) table, col.26 example 12 -----	20-29
X	US 3 718 472 A (OLIVER G ET AL) 27 February 1973 (1973-02-27)	20-24
Y	column 6, line 1 - line 10; claims 1,8; example 5 -----	25-37
A	US 4 617 373 A (PRUETT WAYNE P ET AL) 14 October 1986 (1986-10-14) column 57 -column 61; table 9 -----	1-12, 20-29
A	DATABASE CAPLUS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; Database accession no. 116:173723 XP002238445 abstract & QI ZHONG ET AL.: "Catalytic synthesis of alpha-cyano, beta-unsaturated sulfones in the presence of organotellurium oxide" CHINESE CHEMICAL LETTERS, vol. 2, no. 9, 1991, pages 683-684, -----	20-29
Y	SCHRÖDER E ET AL: "Arzneimittelchemie passage" ARZNEIMITTELCHÉMIE GRUNDLAGEN NERVEN, MUSKELN UND GEWEBE, XX, XX, 1976, pages 30-33, XP002186820 page 32 -----	25-37

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 17-19 and 35-37 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of these compositions.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 03/00032

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0179158	A	25-10-2001	AU 4820101 A	30-10-2001
			WO 0179158 A2	25-10-2001
			EP 1272457 A2	08-01-2003
			GB 2378948 A	26-02-2003
US 5789427	A	04-08-1998	US 5773476 A	30-06-1998
			US 2002068687 A1	06-06-2002
			AU 2096895 A	25-09-1995
			WO 9524190 A2	14-09-1995
US 4632895	A	30-12-1986	CA 1264594 A1	23-01-1990
			DE 3569302 D1	11-05-1989
			EP 0175504 A2	26-03-1986
			JP 1823222 C	10-02-1994
			JP 5033779 B	20-05-1993
			JP 61069064 A	09-04-1986
			CA 1263048 A1	21-11-1989
US 3718472	A	27-02-1973	BE 780011 A1	16-06-1972
US 4617373	A	14-10-1986	AU 579923 B2	15-12-1988
			AU 5513686 A	10-09-1986
			CA 1282528 A1	02-04-1991
			CN 86101649 A , B	21-01-1987
			DE 3667990 D1	08-02-1990
			EP 0215054 A1	25-03-1987
			JP 7116281 B	13-12-1995
			JP 62501856 T	23-07-1987
			KR 9104880 B1	15-07-1991
			TR 22552 A	22-10-1987
			WO 8604904 A1	28-08-1986
			AU 579233 B2	17-11-1988
			AU 5544986 A	10-09-1986
			CA 1261532 A1	26-09-1989
			CN 86101561 A , B	25-02-1987
			DE 3670852 D1	07-06-1990
			EP 0217823 A1	15-04-1987
			ES 8707994 A1	16-11-1987
			ES 8707995 A1	16-11-1987
			JP 7033436 B	12-04-1995
			JP 62501857 T	23-07-1987
			KR 9104934 B1	18-07-1991
			TR 22945 A	08-12-1988
			WO 8604903 A1	28-08-1986
			US 4617374 A	14-10-1986
			ZA 8601129 A	29-10-1986
			ZA 8601130 A	24-09-1986

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.